

ON MANIN'S CONJECTURE FOR SINGULAR DEL PEZZO SURFACES OF DEGREE FOUR, II

R. DE LA BRETÈCHE AND T.D. BROWNING

ABSTRACT. This paper establishes the Manin conjecture for a certain non-split singular del Pezzo surface of degree four $X \subset \mathbb{P}^4$. In fact, if $U \subset X$ is the open subset formed by deleting the lines from X , and H is the usual projective height function on $\mathbb{P}^4(\mathbb{Q})$, then the height zeta function $\sum_{x \in U(\mathbb{Q})} H(x)^{-s}$ is analytically continued to the half-plane $\Re(s) > 17/20$.

1. INTRODUCTION

Let $X \subset \mathbb{P}^4$ be a singular del Pezzo surface of degree four such that $X(\mathbb{Q})$ is Zariski dense in X , and let $U \subset X$ denote the open subset formed by deleting the lines from X . The purpose of this paper is to extend our previous investigation [2] into the asymptotic distribution of rational points on U . For any $x = [x_0, \dots, x_4] \in \mathbb{P}^4(\mathbb{Q})$ such that $x_0, \dots, x_4 \in \mathbb{Z}$ and $\gcd(x_0, \dots, x_4) = 1$, let $H(x) = \max_{0 \leq i \leq 4} |x_i|$ denote the usual anticanonical height function. Then the behaviour of the associated counting function

$$N_{U,H}(B) = \#\{x \in U(\mathbb{Q}) : H(x) \leq B\},$$

as $B \rightarrow \infty$, is predicted by the Manin conjecture [7]. Let $\tilde{X}$ denote the minimal desingularisation of X and let ρ denote the rank of the Picard group $\text{Pic } \tilde{X}$ of $\tilde{X}$. There is a strong version of this conjecture that predicts the existence of a constant $c_{X,H} > 0$, and a monic polynomial $P \in \mathbb{R}[t]$ of degree $\rho - 1$, such that

$$N_{U,H}(B) = c_{X,H} B P(\log B) + O(B^{1-\delta}), \quad (1.1)$$

for some $\delta > 0$. The constant $c_{X,H}$ has received a conjectural interpretation at the hands of Peyre [14], and will be discussed in §2 below. As yet there appears to be no conjectural understanding of the lower order coefficients in this asymptotic formula. The true nature of the error term has been investigated by Swinnerton-Dyer [15], in the setting of diagonal cubic surfaces.

A classification of singular quartic del Pezzo surfaces can be found in the work of Hodge and Pedoe [11, Book IV, §XIII.11]. This shows that up to isomorphism over $\overline{\mathbb{Q}}$, there are 15 possible singularity types that can occur. Coray and Tsfasman [5, Proposition 6.1] have calculated the extended Dynkin diagrams for each type. Given this finite list of surfaces, it is natural to try and develop an arsenal of tools and techniques that permit us to verify the conjectured asymptotic formula (1.1) for each surface on the list. One approach to establishing the Manin conjecture involves studying the height zeta function

$$Z_{U,H}(s) = \sum_{x \in U(\mathbb{Q})} \frac{1}{H(x)^s}.$$

This is defined when $\Re(s)$ is sufficiently large. Once one has proved suitably strong statements about the analytic properties of $Z_{U,H}(s)$, one automatically obtains information about the asymptotic behaviour of $N_{U,H}(B)$ via standard Tauberian

arguments. This approach was present in our previous work [2], where an extensive study was made of the quartic del Pezzo surface

$$x_0x_1 - x_2^2 = x_0x_4 - x_1x_2 + x_3^2 = 0. \quad (1.2)$$

This surface is split over $\mathbb{Q}$ and has a unique singular point, which is of type $\mathbf{D}_5$. In particular, the Picard group of the minimal desingularisation of (1.2) has maximal rank 6. In addition to providing an analytic continuation of the corresponding height zeta function to the half-plane $\Re(s) > 9/10$, an estimate of the shape (1.1) was obtained for any $\delta \in (0, 1/12)$.

The primary goal of this paper is to determine whether the techniques that were developed in [2] can be brought to bear upon a surface that is not split over the ground field $\mathbb{Q}$. Let $X \subset \mathbb{P}^4$ be the surface

$$x_0x_1 - x_2^2 = x_0^2 - x_1x_4 + x_3^2 = 0. \quad (1.3)$$

Then X has a unique singular point $\xi = [0, 0, 0, 0, 1]$, which is of type $\mathbf{D}_4$. In fact X has singularity type $\mathbf{C}_3$ over $\mathbb{Q}$, in the sense of Lipman [12, §24], which becomes a $\mathbf{D}_4$ singularity over $\overline{\mathbb{Q}}$. It is easy to see that the only two lines that are contained in X are

$$\ell_1 : x_1 = x_2 = x_0 - ix_3 = 0, \quad \ell_2 : x_1 = x_2 = x_0 + ix_3 = 0.$$

Clearly both ℓ_1 and ℓ_2 pass through ξ , which is actually the only rational point lying on either line. In particular $N_{U,H}(B) = N_{X,H}(B) + O(1)$. In our previous work [2] the universal torsor was a fundamental ingredient in the resolution of the Manin conjecture for (1.2), much in keeping with the general philosophy. One of the most novel features of our present investigation is that we will be able to establish the Manin conjecture for (1.3) using a certain sub-torsor of the universal torsor.

Our first result concerns the analytic properties of the associated height zeta function $Z_{U,H}(s)$. For any positive integer n , let

$$\chi(n) = \begin{cases} +1 & \text{if } n \equiv 1 \pmod{4}, \\ -1 & \text{if } n \equiv 3 \pmod{4}, \\ 0 & \text{otherwise,} \end{cases}$$

be the real non-principal character modulo 4. Then for $\Re(s) > 0$ we introduce the functions

$$E_1(s+1) = \zeta(2s+1)^2 \zeta(3s+1) \zeta(4s+1) L(2s+1, \chi) L(3s+1, \chi), \quad (1.4)$$

$$E_2(s+1) = \frac{\zeta(9s+3) L(9s+3, \chi)}{\zeta(5s+2)^2 \zeta(6s+2)^2 L(5s+2, \chi) L(6s+2, \chi)^2}. \quad (1.5)$$

It is easily seen that $E_1(s)$ has a meromorphic analytic continuation to the entire complex plane with a single pole at $s = 1$. Similarly it is clear that $E_2(s)$ is holomorphic and bounded on the half-plane $\{s \in \mathbb{C} : \Re(s) \geq 5/6 + \varepsilon\}$, for any $\varepsilon > 0$. For any $\alpha \in \mathbb{R}$ let

$$\mathcal{H}_\alpha = \{s \in \mathbb{C} : \Re(s) \geq \alpha + \varepsilon\}.$$

We are now ready to state our main result.

Theorem 1. *Let $\varepsilon > 0$. Then there exists a constant $\beta \in \mathbb{R}$, and functions $G_1(s), G_2(s)$ that are holomorphic on the half-planes $\mathcal{H}_{3/4}$ and $\mathcal{H}_{17/20}$, respectively, such that for $\Re(s) > 1$ we have*

$$Z_{U,H}(s) = E_1(s)E_2(s)G_1(s) + \frac{12/\pi^2 + 4\beta}{s-1} + G_2(s).$$

In particular $(s-1)^4 Z_{U,H}(s)$ has a holomorphic analytic continuation to the half-plane $\mathcal{H}_{17/20}$. The function $G_1(s)$ is bounded on the half-plane $\mathcal{H}_{3/4}$ and satisfies $G_1(1) \neq 0$, and the function $G_2(s)$ satisfies

$$G_2(s) \ll_\varepsilon (1 + |\Im m(s)|)^{20 \max\{1 - \Re e(s), 0\} / 3 + \varepsilon}$$

on the domain $\mathcal{H}_{17/20}$.

The main step in the proof of Theorem 1 consists of establishing a preliminary estimate for $N_{U,H}(B)$. This will be the object of §§3–4. In §5 this estimate will then be used to deduce the analytic properties of $Z_{U,H}(s)$ presented above. Explicit expressions for β and G_1 can be found in (4.18) and (5.2), respectively. It is interesting to compare Theorem 1 with the corresponding result in our previous work [2, Theorem 1]. It is no surprise that the structure of the two height zeta functions is very similar. Thus in both expressions we have a first term $E_1(s)E_2(s)G_1(s)$ that corresponds to the main term in our preliminary estimate for the counting function, a term $\frac{12}{\pi^2}(s-1)^{-1}$ that corresponds to an isolated conic contained in the surface, and a further “ β -term” involving a constant β that arises through the error in approximating certain arithmetic quantities by real-valued continuous functions. This β -term is one of the most mysterious aspects of our work, and it is interesting to highlight the difference in nature between the constant that appears in Theorem 1 and the corresponding constant obtained in [2, Theorem 1]. Thus whereas the β -term in the latter work relies upon results concerning the equidistribution of squares in a fixed residue class, the β -term in Theorem 1 merely arises through a routine application of integration by parts.

We have already mentioned that the main step in the proof of Theorem 1 involves producing a preliminary estimate for $N_{U,H}(B)$. In §6 we will show how Perron’s formula can be combined with Theorem 1 to extract the following asymptotic formula for $N_{U,H}(B)$.

Theorem 2. *Let $\delta \in (0, 3/32)$. Then there exists a polynomial P of degree 3 such that for any $B \geq 1$ we have*

$$N_{U,H}(B) = BP(\log B) + O(B^{1-\delta}).$$

Moreover, the leading coefficient of P is equal to

$$\frac{\pi^2}{576} \int_0^1 \frac{u^{1/4} du}{\sqrt{1-u}} \prod_p \left(1 - \frac{1}{p}\right)^4 \left(1 - \frac{\chi(p)}{p}\right)^2 \left(1 + \frac{3 + 2\chi(p) + \chi^2(p)}{p} + \frac{\chi^2(p)}{p^2}\right).$$

We will verify in §2 that Theorem 2 is in accordance with Manin’s conjecture. A crucial step in the proof of Theorems 1 and 2 is a bijection that we establish between the rational points on U and the points $(v_1, v_2, y_0, \dots, y_4) \in \mathbb{Z}^7$ such that

$$y_0^4 y_2^2 - v_2 y_1^2 y_4 + y_3^2 = 0. \quad (1.6)$$

Note that v_1 does not appear explicitly in the equation. This step is achieved in §3 via an elementary analysis of the equations defining X . As we have already indicated it is interesting to note that this equation is not an affine embedding of the universal torsor over the minimal desingularisation $\tilde{X}$ of X . Instead it turns out that (1.6) corresponds to a certain sub-torsor of the universal torsor, which reflects the fact that X does not split over the ground field. Theorem 2 seems to signify the first time that the full Manin conjecture has been established without recourse to the universal torsor.

Over the last decade or so the Manin conjecture has been established for a variety of special cases, and it is important to place our investigation in the context of other work. We will say nothing about the situation for non-singular del Pezzo surfaces, or singular del Pezzo surfaces of degree not equal to four. A discussion of

these surfaces can be found in the second author's survey [3]. It turns out that the Manin conjecture has already been established for several singular quartic del Pezzo surfaces by virtue of the fact that the surface is toric, for which there is the general work of Batyrev and Tschinkel [1], or the surface is an equivariant compactification of $\mathbb{G}_a^2$, for which there is the work of Chambert-Loir and Tschinkel [4]. The surface (1.2) studied in [2] falls into this latter category. That (1.3) is not an equivariant compactification of $\mathbb{G}_a^2$ can be seen by mimicking the argument used by Hassett and Tschinkel [10, Remark 4.3] in their analysis of a certain cubic surface. The authors have recently learnt of work due to Derenthal and Tschinkel [6], in which the Manin conjecture is established for the surface

$$x_0x_3 - x_1x_4 = x_0x_1 + x_1x_3 + x_2^2 = 0.$$

This is the split del Pezzo surface of degree four, with singularity type $\mathbf{D}_4$. Their asymptotic formula is weaker than ours, and does not lead to an analytic continuation of the corresponding height zeta function.

Acknowledgements. The authors are grateful to Ulrich Derenthal and Brendan Hassett for several useful conversations relating to universal torsors for singular del Pezzo surfaces. Special thanks are due to Roger Heath-Brown whose ideas led us to the proof of Lemma 6. The paper was finalised while the first author was at the École Normale Supérieure, and the second author was at Oxford University supported by EPSRC grant number GR/R93155/01. The hospitality and financial support of these institutions is gratefully acknowledged. Finally, the authors would like to thank the anonymous referee for his careful reading of the manuscript and numerous useful suggestions.

2. CONFORMITY WITH THE MANIN CONJECTURE

In this section we will review some of the geometry of the surface $X \subset \mathbb{P}^4$, as defined by the pair of quadratic forms

$$Q_1(\mathbf{x}) = x_0x_1 - x_2^2, \quad Q_2(\mathbf{x}) = x_0^2 - x_1x_4 + x_3^2, \quad (2.1)$$

where $\mathbf{x} = (x_0, x_1, x_2, x_3, x_4)$. In particular we will show that Theorem 2 agrees with the Manin conjecture.

Let $\tilde{X}$ denote the minimal desingularisation of X , and let $\pi : \tilde{X} \rightarrow X$ denote the corresponding blow-up map. We let L_i denote the strict transform of the line ℓ_i for $i = 1, 2$, and let $E_1, \dots, E_4$ denote the exceptional curves of π . Then the divisors $E_1, \dots, E_4, L_1, L_2$ satisfy the Dynkin diagram

$$\begin{array}{ccccccc} & & & E_2 & & & \\ & & & | & & & \\ L_1 & \text{---} & E_3 & \text{---} & E_1 & \text{---} & E_4 \text{---} L_2 \end{array}$$

after a possible relabelling of indices. From this it is possible to write down the 6×6 intersection matrix

	E_1	E_2	E_3	E_4	L_1	L_2
E_1	−2	1	1	1	0	0
E_2	1	−2	0	0	0	0
E_3	1	0	−2	0	1	0
E_4	1	0	0	−2	0	1
L_1	0	0	1	0	−1	0
L_2	0	0	0	1	0	−1

which implies that the geometric Picard group $\text{Pic}_{\overline{\mathbb{Q}}} \tilde{X} = \text{Pic}(\tilde{X} \otimes_{\mathbb{Q}} \overline{\mathbb{Q}})$ is generated by $E_1, E_2, E_3, E_4, L_1, L_2$. Moreover the adjunction formula implies that

$$-K_{\tilde{X}} = 4E_1 + 2E_2 + 3(E_3 + E_4) + 2(L_1 + L_2), \quad (2.2)$$

where $-K_{\tilde{X}}$ denotes the anticanonical divisor of $\tilde{X}$. Now if Γ denotes the Galois group of $\mathbb{Q}(i)/\mathbb{Q}$, then it is clear that $\{L_1, L_2\}^\sigma = \{L_1, L_2\}$ for any $\sigma \in \Gamma$. Furthermore, it emerges during the calculation of $\tilde{X}$ that

$$E_1^\sigma = E_1, \quad E_2^\sigma = E_2, \quad \{E_3, E_4\}^\sigma = \{E_3, E_4\},$$

for any $\sigma \in \Gamma$. The Picard group $\text{Pic} \tilde{X}$ of $\tilde{X}$ is therefore the free abelian group generated by $E_1, E_2, E_3 + E_4, L_1 + L_2$. In particular $\rho = 4$ in (1.1), which agrees with Theorem 2.

It remains to discuss the conjectured value of the constant $c_{X,H}$ in (1.1). For this we will follow the presentation adopted in our previous investigation [2, §2], and so we will permit ourselves to be brief. In the notation found there, we see that the conjectured value of the constant in (1.1) is

$$c_{X,H} = \alpha(\tilde{X})\beta(\tilde{X})\tau_H(\tilde{X}). \quad (2.3)$$

Now it follows from [5, Theorem 7.2] that $H^1(\mathbb{Q}, \text{Pic}_{\overline{\mathbb{Q}}} \tilde{X}) = 0$, whence $\beta(\tilde{X}) = 1$. Turning to the value of $\alpha(\tilde{X})$, we have already seen how $-K_{\tilde{X}}$ can be written in terms of the basis for $\text{Pic} \tilde{X}$. Moreover it is easy to check that the cone of effective divisors $\Lambda_{\text{eff}}(\tilde{X}) \subset (\text{Pic} \tilde{X}) \otimes_{\mathbb{Z}} \mathbb{R}$ is also generated by the basis elements of $\text{Pic} \tilde{X}$. This allows us to conclude that

$$\begin{aligned} \alpha(\tilde{X}) &= \text{meas} \{(t_1, t_2, t_3, t_4) \in \mathbb{R}_{\geq 0}^4 : 4t_1 + 2t_2 + 3t_3 + 2t_4 = 1\} \\ &= \frac{1}{2} \text{meas} \{(t_1, t_2, t_3) \in \mathbb{R}_{\geq 0}^3 : 4t_1 + 2t_2 + 3t_3 \leq 1\} \\ &= \frac{1}{288}. \end{aligned} \quad (2.4)$$

The calculation of $\tau_H(\tilde{X})$ is a little more involved, and will be carried out in the following result.

Lemma 1. *We have $\tau_H(\tilde{X}) = \pi^2 \omega_\infty \tau / 16$, where*

$$\omega_\infty = 8 \int_0^1 \frac{u^{1/4} du}{\sqrt{1-u}} \quad (2.5)$$

and

$$\tau = \prod_p \left(1 - \frac{1}{p}\right)^4 \left(1 - \frac{\chi(p)}{p}\right)^2 \left(1 + \frac{3 + 2\chi(p) + \chi^2(p)}{p} + \frac{\chi^2(p)}{p^2}\right). \quad (2.6)$$

Proof. Write $L_p(s, \text{Pic}_{\overline{\mathbb{Q}}} \tilde{X})$ for the local factors of $L(s, \text{Pic}_{\overline{\mathbb{Q}}} \tilde{X})$. Furthermore, let ω_∞ denote the archimedean density of points on X , and let ω_p denote the usual p -adic density of points on X , for any prime p . Then the Tamagawa measure is given by

$$\tau_H(\tilde{X}) = \lim_{s \rightarrow 1} ((s-1)^\rho L(s, \text{Pic}_{\overline{\mathbb{Q}}} \tilde{X})) \omega_\infty \prod_p \frac{\omega_p}{L_p(1, \text{Pic}_{\overline{\mathbb{Q}}} \tilde{X})},$$

where $\rho = 4$ is the rank of $\text{Pic} \tilde{X}$. Our first step is to note that

$$L(s, \text{Pic}_{\overline{\mathbb{Q}}} \tilde{X}) = \zeta(s)^2 \zeta_{\mathbb{Q}(i)}(s)^2.$$

Since $\zeta_{\mathbb{Q}(i)}(s) = \zeta(s)L(s, \chi)$, it then follows that

$$\lim_{s \rightarrow 1} ((s-1)^\rho L(s, \text{Pic}_{\overline{\mathbb{Q}}} \tilde{X})) = L(1, \chi)^2 = \frac{\pi^2}{16}.$$

Furthermore, we plainly have

$$L_p(1, \text{Pic}_{\mathbb{Q}} \tilde{X})^{-1} = \left(1 - \frac{1}{p}\right)^4 \left(1 - \frac{\chi(p)}{p}\right)^2, \quad (2.7)$$

for any prime p .

We proceed by calculating the value of the archimedean density ω_∞ . Let $\|\mathbf{x}\|$ denote the norm $\max_{0 \leq i \leq 4} |x_i|$ for any $\mathbf{x} \in \mathbb{R}^5$. We will follow the method given by Peyre [14] to compute ω_∞ . It will be convenient to parametrise the points via the choice of variables x_0, x_1, x_4 , for which we first observe that

$$\det \begin{pmatrix} \frac{\partial Q_1}{\partial x_2} & \frac{\partial Q_2}{\partial x_2} \\ \frac{\partial Q_1}{\partial x_3} & \frac{\partial Q_2}{\partial x_3} \end{pmatrix} = -4x_2x_3.$$

Now in any real solution to the pair of equations $Q_1(\mathbf{x}) = Q_2(\mathbf{x}) = 0$, the components x_0, x_1 and x_4 must necessarily all share the same sign. Taking into account the fact that $\mathbf{x}$ and $-\mathbf{x}$ represent the same point in $\mathbb{P}^4$, the archimedean density of points on X is therefore equal to

$$\omega_\infty = 4 \int_{\{\mathbf{x} \in \mathbb{R}_{>0}^5 : Q_1(\mathbf{x}) = Q_2(\mathbf{x}) = 0, \|\mathbf{x}\| \leq 1\}} \omega_L(\mathbf{x}),$$

where $\omega_L(\mathbf{x})$ is the Leray form $(4x_2x_3)^{-1} dx_0 dx_1 dx_4$. It follows that

$$\begin{aligned} \omega_\infty &= \int \int \int_{\left\{ \begin{array}{l} x_0, x_1, x_4 \in \mathbb{R}_{>0} : \\ x_1, x_4 \leq 1, x_0^2 < x_1 x_4 \end{array} \right\}} \frac{dx_0 dx_1 dx_4}{\sqrt{x_0 x_1 (x_1 x_4 - x_0^2)}} \\ &= 2 \int \int_{\left\{ \begin{array}{l} x_0, x_1 \in \mathbb{R}_{>0} : \\ x_1 \leq 1, x_0^2 < x_1 \end{array} \right\}} \frac{\sqrt{x_1 - x_0^2}}{x_0^{1/2} x_1^{3/2}} dx_0 dx_1. \end{aligned}$$

The change of variables $u = x_0^2/x_1$ therefore yields

$$\omega_\infty = \int_0^1 \frac{\sqrt{1-u}}{u^{3/4}} du \int_0^1 \frac{dx_1}{x_1^{3/4}} = 8 \int_0^1 \frac{u^{1/4} du}{\sqrt{1-u}},$$

where we have carried out integration by parts to get the last equality. This establishes (2.5).

It remains to calculate the value of $\omega_p = \lim_{r \rightarrow \infty} p^{-3r} N(p^r)$, for any prime p , where

$$N(p^r) = \#\{\mathbf{x} \pmod{p^r} : Q_1(\mathbf{x}) \equiv Q_2(\mathbf{x}) \equiv 0 \pmod{p^r}\}.$$

Although this amounts to a routine calculation, the arguments needed to handle our non-split surface are slightly more subtle, and we have decided to present them in full. To begin with we write $x_0 = p^{k_0} x'_0$ and $x_1 = p^{k_1} x'_1$, with $p \nmid x'_0 x'_1$. Now we have $p^r \mid x_2^2$ if and only if $k_0 + k_1 \geq r$, and there are at most $p^{r/2}$ square roots of zero modulo p^r . When $k_0 + k_1 < r$, it follows that $k_0 + k_1$ must be even and we may write $x_2 = p^{(k_0+k_1)/2} x'_2$, with $p \nmid x'_2$ and

$$x'_0 x'_1 - x'^2_2 \equiv 0 \pmod{p^{r-k_0-k_1}}.$$

The number of possible choices for x'_0, x'_1, x'_2 is therefore

$$h_p(r, k_0, k_1) = \begin{cases} \phi(p^{r-k_0}) \phi(p^{r-(k_0+k_1)/2}) p^{k_0} & \text{if } k_0 + k_1 < r, \\ O(p^{5r/2-k_0-k_1}) & \text{if } k_0 + k_1 \geq r. \end{cases}$$

It remains to determine the number of solutions x_3, x_4 modulo p^r such that

$$p^{2k_0} x'^2_0 - p^{k_1} x'_1 x_4 + x^2_3 \equiv 0 \pmod{p^r}. \quad (2.8)$$

In order to do so we distinguish between four basic cases: either $k_0 + k_1 < r$ and $2k_0 = k_1$, or $k_0 + k_1 < r$ and $2k_0 < k_1$, or $k_0 + k_1 < r$ and $2k_0 > k_1$, or else

$k_0 + k_1 \geq r$. For the first three of these cases we must take care only to sum over values of k_0, k_1 such that $k_0 + k_1$ is even. We will denote by $N_i(p^r)$ the contribution to $N(p^r)$ from the i th case, for $1 \leq i \leq 4$, so that

$$N(p^r) = N_1(p^r) + N_2(p^r) + N_3(p^r) + N_4(p^r). \quad (2.9)$$

We begin by calculating the value of $N_1(p^r)$. For this we write $x_3 = p^{k_3} x'_3$, with $k_3 = \min\{r/2, k_0\} = k_0$. The number of possibilities for x'_3 is p^{r-k_0} , each one leading to precisely p^{2k_0} possible choices for x_4 via (2.8). On noting that k_1 is even, so that k_0 must be even, we may write $k_0 = 2k'_0$, for $0 \leq k'_0 < r/6$. In this way we deduce that

$$N_1(p^r) = \sum_{0 \leq k'_0 < r/6} p^{r+2k'_0} h_p(r, 2k'_0, 4k'_0) = p^{3r}(1 - 1/p) + O(p^{3r-r/6-1}).$$

Next we calculate $N_2(p^r)$. As above we write $x_3 = p^{k_0} x'_3$, and consider the resulting congruence

$$x_0'^2 + x_3'^2 \equiv p^{k_1-2k_0} x'_1 x_4 \pmod{p^{r-2k_0}}.$$

Suppose first that p is odd. Then modulo $p^{k_1-2k_0}$, there are $1 + \chi(p)$ choices for x'_3 , whence there are $(1 + \chi(p))p^{r+k_0-k_1}$ possibilities for x'_3 overall. But then x_4 satisfies a congruence modulo p^{r-k_1} , and there are therefore p^{k_1} ways of choosing x_4 . On summing these contributions over all the relevant values of k_0, k_1 , we obtain

$$\begin{aligned} N_2(p^r) &= \sum_{\substack{k_0+k_1 < r, \ k_0, k_1 \geq 0 \\ 2k_0 < k_1, \ 2|(k_0+k_1)}} (1 + \chi(p)) p^{r+k_0} h_p(r, k_0, k_1) \\ &= 2(1 + \chi(p)) p^{3r-1} (1 + o(1)), \end{aligned}$$

when p is odd. When $p = 2$ the only difference in this calculation is that we must restrict ourselves to the case $k_1 - 2k_0 = 1$, since there are no solutions to the congruence $x_0'^2 + x_3'^2 \equiv 0 \pmod{2^\nu}$ for $\nu \geq 2$. We therefore obtain

$$\begin{aligned} N_2(2^r) &= \sum_{\substack{k_0+k_1 < r, \ k_0, k_1 \geq 0 \\ k_1=2k_0+1, \ 2|(k_0+k_1)}} (1 + \chi(2)) 2^{r+k_0} h_2(r, k_0, k_1) \\ &= 2^{3r-2} (1 + o(1)). \end{aligned}$$

We proceed by calculating $N_3(p^r)$, for which we now write $x_3 = p^{k_3} x'_3$ with $k_3 = \min\{r/2, \lceil k_1/2 \rceil\} = \lceil k_1/2 \rceil$. Here, as throughout this work, $\lceil \alpha \rceil$ denotes the ceiling function for any $\alpha \in \mathbb{R}$, whereas $[\alpha]$ will always denote the integer part of α . With this in mind (2.8) becomes

$$p^{2k_0-k_1} x_0'^2 + p^{2\lceil k_1/2 \rceil - k_1} x_3'^2 \equiv x'_1 x_4 \pmod{p^{r-k_1}}.$$

If k_1 is even, the number of choices for x_3 is $p^{r-k_1/2}$ and this leads to p^{k_1} choices for x_4 . Thus there are $p^{r+k_1/2}$ possibilities for x_3, x_4 if k_1 is even. If k_1 is odd, then p divides x_4 and we find that there are $p^{r+(k_1-1)/2}$ possibilities for x_3 and x_4 . Summing over the relevant values of k_0 and k_1 we therefore obtain

$$\begin{aligned} N_3(p^r) &= \sum_{\substack{k_0+k_1 < r, \ k_0, k_1 \geq 0 \\ 2k_0 > k_1, \ 2|(k_0+k_1)}} p^{r+\lceil k_1/2 \rceil} h_p(r, k_0, k_1) \\ &= p^{3r-2} (p-1)^2 \sum_{\substack{k_0+k_1 < r, \ k_0, k_1 \geq 0 \\ 2k_0 > k_1, \ 2|(k_0+k_1)}} p^{\lceil k_1/2 \rceil - k_1/2 - k_0/2}. \end{aligned}$$

On splitting the summation into four different cases according to the value of k_1 modulo 4, a routine calculation therefore yields

$$\begin{aligned} \frac{N_3(p^r)}{p^{3r-2}(p-1)^2} &= \sum_{\substack{k_0=2k'_0, \ k_1=4k'_1 \\ k'_0 \geq k'_1+1}} p^{-k'_0} + \sum_{\substack{k_0=2k'_0, \ k_1=4k'_1+2 \\ k'_0 \geq k'_1+1}} p^{-k'_0} \\ &+ \sum_{\substack{k_0=2k'_0+1, \ k_1=4k'_1+1 \\ k'_0 \geq k'_1}} p^{-k'_0-1} + \sum_{\substack{k_0=2k'_0+1, \ k_1=4k'_1+3 \\ k'_0 \geq k'_1+1}} p^{-k'_0-1} \\ &= \frac{3p+1}{(p-1)^2} (1 + o(1)), \end{aligned}$$

as $r \rightarrow \infty$.

Finally we calculate the value of $N_4(p^r)$. In this case a straightforward calculation shows that there are at most $2p^{r-k_1+\min\{k_0, k_1/2\}}$ possibilities for x_3 , each one leading to at most p^{k_1} choices for x_4 . We therefore deduce that

$$N_4(p^r) \ll \sum_{k_0+k_1 \geq r} p^{r+\min\{k_0, k_1/2\}} h_p(r, k_0, k_1) \ll p^{3r-r/6}.$$

We may now combine our estimates for $N_1(p^r), \dots, N_4(p^r)$ into (2.9). When p is odd we therefore deduce that

$$\omega_p = \lim_{r \rightarrow \infty} p^{-3r} N(p^r) = 1 + \frac{4 + 2\chi(p)}{p} + \frac{1}{p^2},$$

whereas when $p = 2$ we obtain

$$\omega_2 = \lim_{r \rightarrow \infty} 2^{-3r} N(2^r) = \frac{5}{2}.$$

On combining this with (2.7), we therefore conclude the proof of (2.6), and so complete the proof of Lemma 1. $\square$

We end this section by combining (2.4) and Lemma 1 in (2.3), in order to deduce that the conjectured value of the constant in (1.1) agrees with the value of the leading coefficient in Theorem 2.

3. PRELIMINARY MANOEUVRES

In this section we will establish an alternative expression for $N_{U,H}(B)$, for which we will follow the presentation of [2, §4]. Let us begin by recalling the notation used there. For any $n \geq 2$ we will let Z^{n+1} denote the set of primitive vectors in $\mathbb{Z}^{n+1}$, and similarly, we let N^{n+1} denote the set of primitive vectors in $\mathbb{N}^{n+1}$. Moreover, let $\mathbb{Z}_*^{n+1}$ (resp. $\mathbb{Z}_*^{n+1}$) denote the set of vectors $\mathbf{v} \in \mathbb{Z}^{n+1}$ (resp. $\mathbf{v} \in \mathbb{Z}^{n+1}$) such that $v_0 \cdots v_n \neq 0$. Finally, we will henceforth follow common convention and allow the small parameter $\varepsilon > 0$ to take different values at different points of the argument.

If $x = [\mathbf{x}] \in X \cap \mathbb{P}^4(\mathbb{Q})$ is represented by the vector $\mathbf{x} \in \mathbb{Z}^5$, then it is easy to see that

$$H(x) = \max\{|x_1|, |x_4|\}.$$

Recall the definition (2.1) of the quadratic forms Q_1 and Q_2 . Then it follows that

$$N_{U,H}(B) = \frac{1}{2} \# \left\{ \mathbf{x} \in \mathbb{Z}^5 : \max\{|x_1|, |x_4|\} \leq B, \ Q_1(\mathbf{x}) = Q_2(\mathbf{x}) = 0 \right\} + O(1),$$

since $\mathbf{x}$ and $-\mathbf{x}$ represent the same point in $\mathbb{P}^4$. Let

$$N(Q_1, Q_2; B) = \# \{ \mathbf{x} \in N^5 : \max\{x_1, x_4\} \leq B, \ Q_1(\mathbf{x}) = Q_2(\mathbf{x}) = 0 \}. \quad (3.1)$$

Then we proceed to establish the following basic result.

Lemma 2. *Let $B \geq 1$. Then we have*

$$N_{U,H}(B) = 4N(Q_1, Q_2; B) + \frac{12}{\pi^2}B + O(B^{2/3}).$$

Proof. Let us consider the contribution to $N_{U,H}(B)$ from vectors $\mathbf{x} \in Z^5$ which contain zero components. We claim that

$$N_{U,H}(B) = \frac{1}{2}\#S(Q_1, Q_2; B) + \frac{12}{\pi^2}B + O(B^{2/3}), \quad (3.2)$$

where

$$S(Q_1, Q_2; B) = \{\mathbf{x} \in Z_*^5 : \max\{|x_1|, |x_4|\} \leq B, Q_1(\mathbf{x}) = Q_2(\mathbf{x}) = 0\}.$$

Suppose that $\mathbf{x} \in Z^5$ is a vector such that

$$x_0 = 0, \quad |x_1|, |x_4| \leq B,$$

and $Q_1(\mathbf{x}) = Q_2(\mathbf{x}) = 0$. Then it immediately follows from the first of these equations that $x_2 = 0$, and the second equation implies that $x_3^2 = x_1x_4$. Now either $x_3 = 0$, in which case there are at most 4 possibilities for $\mathbf{x}$, or else we have $\mathbf{x} = \pm(0, a^2, 0, \pm ab, b^2)$ for coprime $a, b \in \mathbb{N}$. Hence the overall contribution from this case is $12B/\pi^2 + O(B^{1/2})$. Suppose now that $\mathbf{x} \in Z^5$ is a vector such that

$$x_1 = 0, \quad |x_4| \leq B,$$

and $Q_1(\mathbf{x}) = Q_2(\mathbf{x}) = 0$. Then a straightforward analysis of these equations reveals that in fact $\mathbf{x} = \pm(0, 0, 0, 0, 1)$. Next we note that any vector with $x_2x_4 = 0$ must have $x_0x_1 = 0$. But such vectors have already been handled above. Finally, if $\mathbf{x} \in Z^5$ satisfies

$$x_3 = 0, \quad |x_1|, |x_4| \leq B,$$

and $Q_1(\mathbf{x}) = Q_2(\mathbf{x}) = 0$, then we must have $x_1x_4 = x_0^2$ and $x_2^2 = x_0x_1$. Since we are only interested in an upper bound it clearly suffices to count non-zero integers x_0, x_2, x_4 , with modulus at most B , such that $\gcd(x_0, x_2, x_4) = 1$ and $x_2^2x_4 = x_0^3$. But then it follows that $(x_0, x_2, x_4) = \pm(a^2b, a^3, b^3)$ for coprime integers a, b , whence the overall contribution is $O(B^{2/3})$. This completes the proof of (3.2).

We now need to relate the cardinality of the set $S(Q_1, Q_2; B)$ to the quantity $N(Q_1, Q_2; B)$. Write $S = S(Q_1, Q_2; B)$ for convenience, and define the four subsets

$$S_{\pm, \pm} = \{\mathbf{x} \in S : \pm x_2 > 0, \pm x_3 > 0\}.$$

Then we clearly have a disjoint union $S = S_{-, -} \cup S_{-, +} \cup S_{+, -} \cup S_{+, +}$, in which each of the four sets has equal size. Hence it follows that $\#S = 4\#S_{+, +}$. Now for any $\mathbf{x} \in S_{+, +}$, we must have that x_0 and x_1 are both positive or both negative, since their product is a square. Similarly x_1 and x_4 both have the same sign, since their product is the sum of two squares. Hence either x_0, x_1, x_4 are all positive, or they are all negative. This therefore establishes the equality

$$\#S_{+, +} = 2N(Q_1, Q_2; B).$$

Upon recalling that $\#S = 4\#S_{+, +}$, and then inserting this into (3.2), we thereby complete the proof of Lemma 2. $\square$

We proceed to equate $N(Q_1, Q_2; B)$ to the number of integral points on a certain affine variety related to X , subject to certain constraints. Let $\mathbf{x} \in N^5$ be any vector counted by $N(Q_1, Q_2; B)$. We begin by considering solutions to the equation $Q_1(\mathbf{x}) = 0$. But it is easy to see that there is a bijection between the set of positive integers x_0, x_1, x_2 such that $x_0x_1 = x_2^2$, and the set of x_0, x_1, x_2 such that

$$x_0 = z_0^2 z_2, \quad x_1 = z_1^2 z_2, \quad x_2 = z_0 z_1 z_2,$$

for $z_0, z_1, z_2 \in \mathbb{N}$, with

$$\gcd(z_0, z_1) = 1. \quad (3.3)$$

We now substitute these values into the equation $Q_2(\mathbf{x}) = 0$, in order to obtain

$$z_0^4 z_2^2 - x_4 z_1^2 z_2 + x_3^2 = 0. \quad (3.4)$$

It is apparent that z_2 divides x_3^2 . Hence we write

$$z_2 = v_2 y_2'^2,$$

for $v_2, y_2' \in \mathbb{N}$ such that v_2 is square-free. It follows that $v_2 y_2'$ divides x_3 , and so there exists $y_3' \in \mathbb{N}$ such that

$$x_3 = v_2 y_2' y_3'.$$

Under these substitutions (3.4) becomes

$$v_2 y_2'^2 z_0^4 - x_4 z_1^2 + v_2 y_3'^2 = 0.$$

At this point it is convenient to deduce a coprimality condition which follows from the assumption made at the outset that $\gcd(x_0, \dots, x_4) = 1$. Recalling the various changes of variables that we have made so far, we easily conclude that

$$\gcd(v_2 y_2', x_4) = 1. \quad (3.5)$$

But then it follows that v_2 must divide z_1^2 in the above equation. Since v_2 is square-free and positive we may conclude that there exists $y_1' \in \mathbb{N}$ such that

$$z_1 = v_2 y_1'.$$

This leads to the equation

$$y_2'^2 z_0^4 - v_2 x_4 y_1'^2 + y_3'^2 = 0, \quad (3.6)$$

in addition to the further coprimality condition

$$\gcd(z_0, v_2 y_1') = 1, \quad (3.7)$$

that follows from (3.3).

Next we let $v_1 = \gcd(y_1', y_3')$. Then $v_1 \in \mathbb{N}$ and we may write

$$z_0 = y_0, \quad y_1' = v_1 y_1, \quad y_3' = v_1 y_3, \quad (3.8)$$

for $y_0, y_1, y_3 \in \mathbb{N}$. Under this change of variables we deduce from (3.7) and the definitions of v_1, v_2 , that

$$|\mu(v_2)| = 1, \quad \gcd(y_1, y_3) = \gcd(y_0, v_1 v_2 y_1) = 1, \quad (3.9)$$

where $\mu(n)$ denotes the Möbius function for any $n \in \mathbb{N}$. We proceed by substituting (3.8) into (3.6). This leads to the equation

$$y_0^4 y_2'^2 - v_1^2 v_2 x_4 y_1^2 + v_1^2 y_3^2 = 0,$$

from which it follows that $v_1^2 \mid y_0^4 y_2'^2$. In view of (3.9) it follows that there exist $y_2, y_4 \in \mathbb{N}$ such that

$$x_4 = y_4, \quad y_2' = v_1 y_2.$$

Our investigation has therefore produced the equation

$$y_0^4 y_2^2 - v_2 y_1^2 y_4 + y_3^2 = 0, \quad (3.10)$$

and (3.5) becomes

$$\gcd(y_4, v_1 v_2 y_2) = 1. \quad (3.11)$$

We take a moment to deduce two further coprimality conditions

$$\gcd(y_2, y_3) = 1, \quad \gcd(y_2, v_2 y_1) = 1. \quad (3.12)$$

Suppose that there exists a prime divisor p common to both y_2 and y_3 . Then it follows from (3.10) that $p^2 \mid v_2 y_1^2 y_4$. But (3.9) and (3.11) together imply that $\gcd(p, y_1 y_4) = 1$. Hence $p^2 \mid v_2$, which is impossible since v_2 is square-free. The

equation (3.10) and the fact that $\gcd(y_2, y_3) = 1$ clearly yield the second relation in (3.12). Combining (3.12) with (3.9) and (3.11), we therefore obtain the relations

$$\gcd(y_0, v_1 v_2 y_1) = \gcd(y_3, y_1 y_2) = \gcd(y_4, v_1 v_2 y_2) = 1, \quad (3.13)$$

and

$$|\mu(v_2)| = 1, \quad \gcd(y_2, v_2 y_1) = 1. \quad (3.14)$$

At this point we may summarise our argument as follows. Let $\mathcal{T}$ denote the set of $(\mathbf{v}, \mathbf{y}) = (v_1, v_2, y_0, \dots, y_4) \in \mathbb{N}^7$ such that (3.10), (3.13) and (3.14) hold. Then for any $\mathbf{x} \in N^5$ counted by $N(Q_1, Q_2; B)$, we have shown that there exists $(\mathbf{v}, \mathbf{y}) \in \mathcal{T}$ such that

$$\begin{aligned} x_0 &= v_1^2 v_2 y_0^2 y_2^2, & x_1 &= v_1^4 v_2^3 y_1^2 y_2^2, & x_2 &= v_1^3 v_2^2 y_0 y_1 y_2^2, \\ x_3 &= v_1^2 v_2 y_2 y_3, & x_4 &= y_4. \end{aligned}$$

Conversely, given any $(\mathbf{v}, \mathbf{y}) \in \mathcal{T}$, the point $\mathbf{x}$ given above will be a solution of the equations $Q_1(\mathbf{x}) = Q_2(\mathbf{x}) = 0$, with $\mathbf{x} \in N^5$. To check that $\gcd(x_0, \dots, x_4) = 1$, we first observe that

$$\gcd(x_0, x_1, x_2) = v_1^2 v_2 y_2^2 \gcd(y_0^2, v_1^2 v_2^2 y_1^2, v_1 v_2 y_0 y_1) = v_1^2 v_2 y_2^2,$$

by (3.13). But then (3.13) also implies that

$$\gcd(x_0, x_1, x_2, x_3) = v_1^2 v_2 y_2 \gcd(y_2, y_3) = v_1^2 v_2 y_2,$$

whence

$$\gcd(x_0, \dots, x_4) = \gcd(v_1^2 v_2 y_2, y_4) = 1.$$

On observing that the inequality $y_4 \leq B$ is equivalent to $y_0^4 y_2^2 + y_3^2 \leq B v_2 y_1^2$, by (3.10), we therefore conclude the proof of the following result.

Lemma 3. *Let $B \geq 1$. Then we have*

$$N(Q_1, Q_2; B) = \#\left\{(\mathbf{v}, \mathbf{y}) \in \mathcal{T} : v_1^4 v_2^3 y_1^2 y_2^2 \leq B, \ y_0^4 y_2^2 + y_3^2 \leq B v_2 y_1^2\right\}.$$

It will become clear in subsequent sections that the equation (3.10) is a pivotal ingredient in our proof of Theorems 1 and 2.

4. THE MAIN OFFENSIVE

In this section we use Lemma 2 and Lemma 3 to obtain an initial estimate for $N_{U,H}(B)$, which will then be used to deduce the statement of Theorem 1 in §5. Before beginning this task, it will be helpful to first outline our strategy. It follows from the statement of Lemma 3 that any vector $(\mathbf{v}, \mathbf{y})$ counted by $N(Q_1, Q_2; B)$ is constrained to lie in the region defined by (3.10) and the inequalities

$$v_1^4 v_2^3 y_1^2 y_2^2 \leq B, \quad y_0^4 y_2^2 + y_3^2 \leq B v_2 y_1^2.$$

The bulk of our work will be taken up with handling the summation over the variables y_0, y_3 and y_4 , for fixed values of $(\mathbf{v}, y_1, y_2)$. The essential trick will be to view (3.10) as a congruence condition in order to handle the summation over y_4 . Then we will need to count values of y_3 such that

$$y_3 \leq \sqrt{B v_2 y_1^2 - y_0^4 y_2^2} = Y_3, \quad (4.1)$$

say, subject to certain coprimality conditions, and then finally values of y_0 such that

$$y_0 \leq \frac{B^{1/4} v_2^{1/4} y_1^{1/2}}{y_2^{1/2}} = Y_0, \quad (4.2)$$

say, subject to certain coprimality conditions. The overall contribution from y_0, y_3 and y_4 , which we henceforth denote by $S(\mathbf{v}, y_1, y_2)$, will be estimated in §4.2. Finally

in §4.3 we will sum $S(\mathbf{v}, y_1, y_2)$ over the remaining values of $\mathbf{v}, y_1, y_2$ subject to the inequality

$$v_1^4 v_2^3 y_1^2 y_2^2 \leq B, \quad (4.3)$$

and certain coprimality conditions.

4.1. Congruences and equidistribution. It will be convenient to collect together some technical facts about congruences that will be needed in §4.2 and §4.3. We begin by discussing the arithmetic function $\eta(q)$, defined to be the number of square roots of -1 modulo q . The Chinese Remainder Theorem implies that $\eta(q)$ is multiplicative, and for any $\nu \geq 1$ we have

$$\eta(p^\nu) = \begin{cases} 2 & \text{if } p \equiv 1 \pmod{4}, \\ 0 & \text{if } p \equiv 3 \pmod{4}, \\ 1 & \text{if } p = 2, \nu = 1, \\ 0 & \text{if } p = 2, \nu \geq 2. \end{cases}$$

Let $\omega(n)$ denote the number of distinct prime factors of $n \in \mathbb{N}$, and recall the definition of χ , the real non-principal character modulo 4. Then it is not difficult to see that we have

$$\eta(q) \leq \sum_{d|q} |\mu(d)| \chi(d) \leq 2^{\omega(q)}, \quad (4.4)$$

for any positive integer q .

Define the real-valued function $\psi(t) = \{t\} - 1/2$, where $\{\alpha\}$ denotes the fractional part of $\alpha \in \mathbb{R}$. Then ψ is periodic with period 1. We proceed by recording the following simple estimate.

Lemma 4. *Let $a, q \in \mathbb{Z}$ be such that $q > 0$, and let $t \geq 0$. Then we have*

$$\#\{0 < n \leq t : n \equiv a \pmod{q}\} = \frac{t}{q} + r(t; a, q),$$

where

$$r(t; a, q) = \psi\left(-\frac{a}{q}\right) - \psi\left(\frac{t-a}{q}\right).$$

Proof. This follows on taking $t_1 = 0$ and $t_2 = t$ in [2, Lemma 3]. $\square$

We will also need to prove a result about the average order of the function ψ , that plays the same role in this work that [2, Lemma 4] did in our previous work. A crucial ingredient in this will be the following Diophantine approximation result, which may be of independent interest.

Lemma 5. *Let $q \in \mathbb{N}$ and let ϱ be a square root of -1 modulo q . For each non-zero integer b , there exist coprime integers u, v such that*

$$\left| \frac{b\varrho}{q} - \frac{u}{v} \right| \leq \frac{1}{v\sqrt{2q}} \quad (4.5)$$

and

$$\frac{\sqrt{q/2}}{|b|} \leq v \leq \sqrt{2q}. \quad (4.6)$$

Proof. By Dirichlet's approximation theorem we may find coprime integers u, v such that $1 \leq v \leq \sqrt{2q}$ and (4.5) holds. We claim that any such v automatically satisfies (4.6), for which it clearly remains to establish the lower bound. To do so we first note that $|r| \leq \sqrt{q/2}$, where r denotes the residue of $b\varrho v$ modulo q . But then it follows that

$$r^2 \equiv b^2 \varrho^2 v^2 \equiv -b^2 v^2 \pmod{q},$$

since ϱ is a square-root of -1 modulo q , whence $q \mid r^2 + b^2 v^2$. Thus we obtain the system of inequalities

$$q \leq r^2 + b^2 v^2 \leq q/2 + b^2 v^2,$$

which thereby gives the lower bound in (4.6). $\square$

Given any real-valued function f defined on an interval I , and given coefficients $c_1, c_2 \in \mathbb{R}$ and $q \in \mathbb{N}$, define the sum

$$S_I(f; c_1, c_2, q) = \sum_{x \in \mathbb{Z} \cap I} \psi\left(\frac{c_1 f(x) - c_2 x^2}{q}\right).$$

We proceed by establishing the following result, which permits $S_I(f; c_1, c_2, q)$ to be estimated over short intervals I under suitable hypotheses.

Lemma 6. *Let $b, q \in \mathbb{Z}$ with $q > 0$ and $b \neq 0$, and let ϱ be a square root of -1 modulo q . Let $\gamma \in \mathbb{R}$. Given a bounded interval $I \subset \mathbb{R}$, let $f : I \rightarrow \mathbb{R}$ be a continuously differentiable function on I . Write*

$$\lambda(f) = \int_I |f'(t)| dt,$$

and define

$$g_{\alpha, \beta}(t) = f(t) - \alpha - \beta t,$$

for any $\alpha, \beta \in \mathbb{R}$. Then for any $H \geq 1$ we have

$$S_I(f; \gamma, b\varrho, q) \ll \frac{\mathfrak{m}(I)}{H} + \log(HR) \left(1 + \frac{|\gamma| H \lambda(g_{\alpha, \beta})}{q}\right) \left(q^{1/4} + \frac{|b|^{1/2} H^{1/2} \mathfrak{m}(I)}{q^{1/4}}\right),$$

where $\mathfrak{m}(I) = \text{meas}(I) + 1$ and R is such that $I \subseteq [-R, R]$.

Proof. We employ Vaaler's trigonometric polynomial approximation [17] to ψ . This implies that for any $H \geq 1$, there exist coefficients c_h such that

$$\psi(t) \leq \sum_{0 < |h| \leq H} c_h e(ht) + O\left(\frac{1}{H}\right),$$

with $c_h \ll 1/|h|$ and $e(\theta) = \exp(2\pi i \theta)$. It therefore follows that

$$S_I(f; \gamma, b\varrho, q) \ll \frac{\mathfrak{m}(I)}{H} + \sum_{0 < |h| \leq H} \frac{1}{|h|} \left| \sum_{x \in \mathbb{Z} \cap I} e\left(\frac{h\gamma f(x) - bh\varrho x^2}{q}\right) \right|, \quad (4.7)$$

for any $H \geq 1$.

Define the sum

$$S_{h,Y}(F) = \sum_{Y/2 < y \leq Y} e\left(\frac{hF(y) - bh\varrho y^2}{q}\right),$$

for fixed values of $h \in \mathbb{N}$, $Y \geq 1$ and any real valued function $F : (Y/2, Y] \rightarrow \mathbb{R}$. We begin by estimating the sum when $F(t) = \alpha + \beta t$. Mimicking the proof of Weyl's inequality, we obtain

$$\begin{aligned} |S_{h,Y}(\alpha + \beta t)|^2 &= \left| \sum_{\substack{Y/2 < y \leq Y \\ Y/2 < y+z \leq Y}} e\left(\frac{-\beta h z + bh\varrho(z^2 + 2yz)}{q}\right) \right| \\ &\leq \sum_{|z| \leq Y} \left| \sum_{y \in I(z)} e\left(\frac{2bh\varrho y z}{q}\right) \right|, \end{aligned}$$

for some interval $I(z)$ of length $O(Y)$, depending on z . For any $\alpha \in \mathbb{R}$, let $\|\alpha\|$ denote the distance from α to the nearest integer. Then it follows that

$$|S_{h,Y}(\alpha + \beta t)|^2 \ll \sum_{|z| \leq Y} \min \left\{ Y, \frac{1}{\|2b\varrho h z/q\|} \right\}.$$

For any $\alpha \in \mathbb{R}$ and any $\delta > 0$, let $T(N; \alpha, \delta)$ denote the number of positive integers $n \leq N$ such that $\|\alpha n\| \leq \delta$. A result due to Heath-Brown [9, Lemma 6] yields

$$T(N; \alpha, \delta) \leq 4(1 + v\delta) \left(1 + \frac{N}{v} \right), \quad (4.8)$$

provided that α has a rational approximation $|\alpha - u/v| \leq 1/v^2$ with coprime integers u, v such that $v \neq 0$. On combining (4.8) with Lemma 5, we therefore deduce that

$$\begin{aligned} |S_{h,Y}(\alpha + \beta t)|^2 &\ll \max_{1/Y \leq \delta \leq 1} \frac{T(Y; 2bh\varrho/q, \delta)}{\delta} \\ &\ll (Y + v) \left(1 + \frac{Y}{v} \right) \\ &\ll \left(q^{1/4} + \frac{|bh|^{1/2}Y}{q^{1/4}} \right)^2, \end{aligned}$$

whence

$$S_{h,Y}(\alpha + \beta t) \ll q^{1/4} + \frac{|bh|^{1/2}Y}{q^{1/4}}.$$

Given any $\gamma \in \mathbb{R}$ and any function $f : I \rightarrow \mathbb{R}$ as in the statement of the lemma, it now follows from an application of partial summation that

$$\begin{aligned} S_{h,Y}(\gamma f) &= S_{h,Y}(\gamma g_{\alpha,\beta}(t) + \alpha' + \beta' t) \\ &\ll \left(1 + \frac{h|\gamma|\lambda(g_{\alpha,\beta})}{q} \right) \left(q^{1/4} + \frac{|bh|^{1/2}Y}{q^{1/4}} \right), \end{aligned}$$

with $\alpha' = \alpha\gamma$ and $\beta' = \beta\gamma$. Finally we employ this estimate in (4.7), and sum over dyadic intervals for $Y \ll R$, in order to complete the proof of Lemma 6. $\square$

4.2. Summation over the variables y_0, y_3 and y_4 . Let $(\mathbf{v}, y_1, y_2) \in \mathbb{N}^4$ satisfy (3.14) and (4.3). As indicated above, we will denote the triple summation over y_0, y_3 and y_4 by $S(\mathbf{v}, y_1, y_2)$. In this summation we will need to take into account the coprimality conditions (3.13). We begin by treating the condition $\gcd(y_4, v_1 v_2) = 1$. Note here that the condition $\gcd(y_4, y_2) = 1$ follows immediately from (3.10) and the remaining conditions. A Möbius inversion yields

$$S(\mathbf{v}, y_1, y_2) = \sum_{k_4 | v_1 v_2} \mu(k_4) S_{k_4},$$

where the definition of S_{k_4} is as for $S(\mathbf{v}, y_1, y_2)$ but with the extra condition that $k_4 \mid y_4$ and without the condition $\gcd(y_4, v_1 v_2 y_2) = 1$. Now it is straightforward to deduce from (3.10), (3.13) and (3.14), that

$$\gcd(k_4, y_2) = \gcd(v_1 v_2, y_2, y_3^2) = 1,$$

for any k_4 dividing $v_1 v_2$ and y_4 . It follows that

$$S(\mathbf{v}, y_1, y_2) = \sum_{\substack{y_0 \leq Y_0 \\ \gcd(y_0, v_1 v_2 y_1) = 1}} \sum_{\substack{k_4 | v_1 v_2 \\ \gcd(k_4, y_2) = 1}} \mu(k_4) S'_{k_4},$$

where

$$S'_{k_4} = \# \{ y_3 \in \mathbb{N} : \gcd(y_3, y_1 y_2) = 1, y_3 \leq Y_3, y_3^2 \equiv -y_0^4 y_2^2 \pmod{k_4 v_2 y_1^2} \},$$

and Y_3 is given by (4.1). Let ϱ be a solution of the congruence

$$\varrho^2 \equiv -1 \pmod{k_4 v_2 y_1^2}.$$

Then a little thought reveals that

$$\gcd(\varrho y_0^2 y_2, k_4 v_2 y_1^2) = 1, \quad (4.9)$$

for any $k_4 \mid v_1 v_2$ such that $\gcd(k_4, y_2) = 1$. We may therefore conclude that

$$S(\mathbf{v}, y_1, y_2) = \sum_{\substack{y_0 \leq Y_0 \\ \gcd(y_0, v_1 v_2 y_1) = 1}} \sum_{\substack{k_4 \mid v_1 v_2 \\ \gcd(k_4, y_2) = 1}} \mu(k_4) \sum_{\substack{\varrho \leq k_4 v_2 y_1^2 \\ \varrho^2 \equiv -1 \pmod{k_4 v_2 y_1^2}}} S_{k_4}(\varrho),$$

where

$$S_{k_4}(\varrho) = \# \{y_3 \in \mathbb{N} : \gcd(y_3, y_2) = 1, y_3 \leq Y_3, y_3 \equiv \varrho y_0^2 y_2 \pmod{k_4 v_2 y_1^2}\}.$$

In order to estimate $S_{k_4}(\varrho)$ we employ a further Möbius inversion to treat the coprimality condition $\gcd(y_3, y_2) = 1$. Thus we have

$$S_{k_4}(\varrho) = \sum_{k_2 \mid y_2} \mu(k_2) S_{k_2, k_4}(\varrho),$$

where

$$S_{k_2, k_4}(\varrho) = \# \{y'_3 \in \mathbb{N} : y'_3 \leq Y_3/k_2, k_2 y'_3 \equiv \varrho y_0^2 y_2 \pmod{k_4 v_2 y_1^2}\}.$$

We must therefore estimate the number of positive integers contained in a certain interval, which belong to a certain arithmetic progression. For fixed values of $\mathbf{v}, y_0, y_1, y_2, k_2, k_4, \varrho$, it follows from (4.9) that there exists a unique positive integer $b \leq k_4 v_2 y_1^2$ such that $\gcd(b, k_4 v_2 y_1^2) = 1$ and

$$b k_2 \equiv \varrho y_2 \pmod{k_4 v_2 y_1^2}. \quad (4.10)$$

We may therefore employ Lemma 4 to deduce that

$$\begin{aligned} S_{k_2, k_4}(\varrho) &= \# \{y'_3 \in \mathbb{N} : y'_3 \leq Y_3/k_2, y'_3 \equiv b y_0^2 \pmod{k_4 v_2 y_1^2}\} \\ &= \frac{Y_3}{k_2 k_4 v_2 y_1^2} + r(Y_3/k_2; b y_0^2, k_4 v_2 y_1^2), \end{aligned}$$

where

$$r(Y_3/k_2; b y_0^2, k_4 v_2 y_1^2) = \psi\left(\frac{-b y_0^2}{k_4 v_2 y_1^2}\right) - \psi\left(\frac{Y_3/k_2 - b y_0^2}{k_4 v_2 y_1^2}\right). \quad (4.11)$$

Putting everything together we have therefore established that

$$S(\mathbf{v}, y_1, y_2) = \vartheta(\mathbf{v}, y_1, y_2) \sum_{\substack{y_0 \leq Y_0 \\ \gcd(y_0, v_1 v_2 y_1) = 1}} \frac{Y_3}{v_2 y_1^2} + R_1(B; \mathbf{v}, y_1, y_2), \quad (4.12)$$

with

$$\begin{aligned} \vartheta(\mathbf{v}, y_1, y_2) &= \sum_{\substack{k_4 \mid v_1 v_2 \\ \gcd(k_4, y_2) = 1}} \frac{\mu(k_4) \eta(k_4 v_2 y_1^2)}{k_4} \sum_{k_2 \mid y_2} \frac{\mu(k_2)}{k_2} \\ &= \frac{\phi(y_2)}{y_2} \sum_{\substack{k_4 \mid v_1 v_2 \\ \gcd(k_4, y_2) = 1}} \frac{\mu(k_4) \eta(k_4 v_2 y_1^2)}{k_4}, \end{aligned}$$

and

$$R_1(B; \mathbf{v}, y_1, y_2) = \sum_{\substack{k_4 \mid v_1 v_2 \\ \gcd(k_4, y_2) = 1}} \mu(k_4) \sum_{\substack{\varrho \leq k_4 v_2 y_1^2 \\ \varrho^2 \equiv -1 \pmod{k_4 v_2 y_1^2}}} \sum_{k_2 \mid y_2} \mu(k_2) F_1(B). \quad (4.13)$$

Here we have set

$$F_1(B) = F_1(B; \mathbf{v}, y_1, y_2, k_2, k_4, \varrho) = \sum_{\substack{y_0 \leq Y_0 \\ \gcd(y_0, v_1 v_2 y_1) = 1}} r(Y_3/k_2; b y_0^2, k_4 v_2 y_1^2),$$

and a straightforward calculation reveals that

$$\vartheta(\mathbf{v}, y_1, y_2) = \frac{\eta(v_2 y_1^2) \phi(y_2)}{y_2} \prod_{\substack{p|v_1 \\ p \nmid v_2 y_1 y_2}} \left(1 - \frac{1 + \chi(p)}{p}\right) \prod_{p|v_2 \gcd(v_1, y_1)} \left(1 - \frac{\chi(p)}{p}\right). \quad (4.14)$$

We have used here the fact that whenever $\eta(v_2 y_1^2) \neq 0$, any prime divisor of $v_2 \gcd(v_1, y_1)$ must be congruent to 1 modulo 4. We proceed by establishing the following result.

Lemma 7. *Let $\varepsilon > 0$. Then for any $B \geq 1$ we have*

$$\sum_{\substack{v_1, v_2, y_1, y_2 \in \mathbb{N} \\ (3.14), (4.3) \text{ hold}}} R_1(B; v_1, v_2, y_1, y_2) \ll_{\varepsilon} B^{17/20+\varepsilon}.$$

Proof. We begin with a Möbius inversion to remove the coprimality condition in the summation over y_0 in the definition of $F_1(B)$. Thus it follows from (4.11) that $F_1(B) = F_1^a(B) - F_1^b(B)$, with

$$F_1^a(B) = \sum_{k_0 | v_1 v_2 y_1} \mu(k_0) \sum_{y_0 \leq Y_0/k_0} \psi\left(\frac{-b k_0^2 y_0^2}{k_4 v_2 y_1^2}\right),$$

and

$$F_1^b(B) = \sum_{k_0 | v_1 v_2 y_1} \mu(k_0) \sum_{y_0 \leq Y_0/k_0} \psi\left(\frac{Y_3/k_2 - b k_0^2 y_0^2}{k_4 v_2 y_1^2}\right).$$

On writing $y_2 = k_2 y_2'$, (4.10) implies that $b \equiv \varrho y_2' \pmod{k_4 v_2 y_1^2}$. Here we recall that ϱ is a square root of -1 modulo $k_4 v_2 y_1^2$. We now set

$$q = k_4 v_2 y_1^2. \quad (4.15)$$

An application of Lemma 6 with $\gamma = 0$ yields

$$\begin{aligned} F_1^a(B) &\ll_{\varepsilon} \sum_{k_0 | v_1 v_2 y_1} B^{\varepsilon} \left(\frac{Y_0}{H} + q^{1/4} + \frac{y_2'^{1/2} H^{1/2} Y_0}{q^{1/4}} \right) \\ &\ll_{\varepsilon} B^{\varepsilon} \left(\frac{Y_0}{H} + (k_4 v_2 y_1^2)^{1/4} + \frac{y_2'^{1/2} H^{1/2} Y_0}{(v_2 y_1^2)^{1/4}} \right), \end{aligned}$$

for any $H \geq 1$. Substituting this into (4.13), we therefore obtain an overall contribution of

$$\ll_{\varepsilon} B^{\varepsilon} \left(\frac{Y_0}{H} + v_1^{1/4} v_2^{1/2} y_1^{1/2} + \frac{y_2'^{1/2} H^{1/2} Y_0}{v_2^{1/4} y_1^{1/2}} \right).$$

to $R_1(B; \mathbf{v}, y_1, y_2)$ from $F_1^a(B)$. Here, we have used the estimate (4.4) for $\eta(k_4 v_2 y_1^2)$. It remains to sum this estimate over all $v_1, v_2, y_1, y_2 \in \mathbb{N}$ satisfying (3.14) and (4.3). Recall the definition (4.2) of Y_0 . Then for any $\varepsilon > 0$ we obtain the overall contribution

$$\begin{aligned} &\ll_{\varepsilon} B^{\varepsilon} \sum_{\substack{v_1, v_2, y_1, y_2 \in \mathbb{N} \\ y_1 \leq B^{1/2}/(v_1^{3/2} v_2)}} \left(\frac{B^{1/4} v_2^{1/4} y_1^{1/2}}{H y_2^{1/2}} + v_1^{1/4} v_2^{1/2} y_1^{1/2} + B^{1/4} H^{1/2} \right) \\ &\ll_{\varepsilon} B^{\varepsilon} \left(\frac{B}{H} + B^{3/4} H^{1/2} \right) \end{aligned}$$

to $\sum_{v_1, v_2, y_1, y_2} R_1(B; \mathbf{v}, y_1, y_2)$ from the term $F_1^a(B)$. This term therefore contributes $O_\varepsilon(B^{5/6+\varepsilon})$ to Lemma 7, on taking $H = B^{1/6}$, which is satisfactory.

We must now turn to the overall contribution from the term $F_1^b(B)$. Define the function $f(t) = \sqrt{1-t^4}$. Then it follows from (4.1) that

$$\frac{Y_3}{k_2} = \sqrt{Bv_2y_1^2}f(y_0/Y'_0),$$

in the above expression for $F_1^b(B)$, where we have set $Y'_0 = Y_0/k_0$ for short. We will estimate $F_1^b(B)$ by breaking the summation over y_0 into intervals of length $O(Y'_0/M)$, for any $M \geq 1$ to be selected in due course. Let

$$U_m = \sum_{mY'_0/M < y_0 \leq (m+1)Y'_0/M} \psi\left(\frac{\sqrt{Bv_2y_1^2}f(y_0/Y'_0) - y_2'k_0^2y_0^2}{q}\right),$$

for each $m \in \mathbb{N}$, where q is given by (4.15). We wish to apply Lemma 6 to estimate U_m , in which we will take $\gamma = \sqrt{Bv_2y_1^2}$.

Let us begin by handling U_m for $m \leq M-2$. Here we take

$$\beta = \frac{f'(m/M)}{M}, \quad \alpha = f(m/M) - \frac{mf'(m/M)}{M^2},$$

and proceed to consider the size of $g_{\alpha,\beta}(t/Y'_0)$ on the range of summation. An application of the mean value theorem implies that for $mY'_0/M < t \leq (m+1)Y'_0/M$ we have

$$\begin{aligned} g_{\alpha,\beta}(t/Y'_0) &= f(t/Y'_0) - f(m/M) - \frac{1}{M}f'(m/M)\left(\frac{t}{Y'_0} - \frac{m}{M}\right) \\ &\ll \frac{1}{M^2} \sup_{\tau \in (m, m+1]} |f''(\tau/M)| \\ &\ll \frac{(1 - (m+1)^4/M^4)^{-3/2}}{M^2}. \end{aligned}$$

On making the change of variables $m = M - m' - 1$, so that

$$(1 - (m+1)^4/M^4)^{-3/2} \ll (M/m')^{3/2},$$

we therefore deduce that

$$\begin{aligned} \sum_{1 \leq m \leq M-2} \lambda(g_{\alpha,\beta}(t/Y'_0)) &\ll M^{-2} \sum_{1 \leq m \leq M-2} (1 - (m+1)^4/M^4)^{-3/2} \\ &\ll M^{-1/2} \sum_{1 \leq m' \leq M-2} m'^{-3/2} \ll M^{-1/2}. \end{aligned}$$

It now follows from Lemma 6 that

$$\sum_{1 \leq m \leq M-2} U_m \ll_\varepsilon B^\varepsilon \left(\frac{Y_0}{H} + \left(M + \frac{B^{1/2}H}{v_2^{1/2}y_1M^{1/2}} \right) \left(v_1^{1/4}v_2^{1/2}y_1^{1/2} + \frac{y_2^{1/2}H^{1/2}Y_0}{v_2^{1/4}y_1^{1/2}M} \right) \right).$$

To estimate U_m for $m = M-1$ or M , we choose $\beta = \alpha = 0$ in Lemma 6. In particular it follows that $g_{\alpha,\beta}(t/Y'_0) = f(t/Y'_0) \ll M^{-1/2}$ and so

$$\lambda(g_{\alpha,\beta}(t/Y'_0)) \ll M^{-1/2},$$

for $mY'_0/M < t \leq (m+1)Y'_0/M$ and $M-2 < m \leq M$. We therefore deduce that

$$\sum_{M-2 < m \leq M} U_m \ll_\varepsilon B^\varepsilon \left(\frac{Y_0}{H} + \left(1 + \frac{B^{1/2}H}{v_2^{1/2}y_1M^{1/2}} \right) \left(v_1^{1/4}v_2^{1/2}y_1^{1/2} + \frac{y_2^{1/2}H^{1/2}Y_0}{v_2^{1/4}y_1^{1/2}M} \right) \right),$$

which once combined with the above leads to the conclusion that

$$\begin{aligned} F_1^b(B) &\ll_\varepsilon B^\varepsilon \left(\frac{Y_0}{H} + \left(M + \frac{B^{1/2}H}{v_2^{1/2}y_1M^{1/2}} \right) \left(v_1^{1/4}v_2^{1/2}y_1^{1/2} + \frac{y_2^{1/2}H^{1/2}Y_0}{v_2^{1/4}y_1^{1/2}M} \right) \right) \\ &\ll_\varepsilon B^\varepsilon \left(\frac{Y_0}{H} + \frac{B^{1/3}H^{2/3}}{v_2^{1/3}y_1^{2/3}} \left(v_1^{1/4}v_2^{1/2}y_1^{1/2} + \frac{y_2^{1/2}Y_0v_2^{1/12}y_1^{1/6}}{B^{1/3}H^{1/6}} \right) \right), \end{aligned}$$

on choosing $M = H^{2/3}B^{1/3}/(v_2^{1/3}y_1^{2/3})$. Note that $M \geq 1$ since $v_2y_1^2 \leq B$. Recall the definition (4.2) of Y_0 and let $\varepsilon > 0$. Then arguing as above, we conclude that we have an overall contribution

$$\ll_\varepsilon B^\varepsilon \sum_{v_1, v_2, y_1, y_2} \left(\frac{B^{1/4}v_2^{1/4}y_1^{1/2}}{Hy_2^{1/2}} + \frac{v_1^{1/4}v_2^{1/6}B^{1/3}H^{2/3}}{y_1^{1/6}} + B^{1/4}H^{1/2} \right)$$

to $\sum_{v_1, v_2, y_1, y_2} R_1(B; \mathbf{v}, y_1, y_2)$ from the term $F_1^b(B)$. Here the summation is over $v_1, v_2, y_1, y_2 \in \mathbb{N}$ such that $y_1 \leq B^{1/2}/(v_1^2v_2^{3/2}y_2)$. Taking $H = y_1^{3/10}$, we therefore obtain the satisfactory contribution

$$\begin{aligned} &\ll_\varepsilon B^\varepsilon \sum_{v_1, v_2, y_1, y_2} \left(\frac{B^{1/4}v_2^{1/4}y_1^{1/5}}{y_2^{1/2}} + v_1^{1/4}v_2^{1/6}y_1^{1/30}B^{1/3} + B^{1/4}y_1^{3/20} \right) \\ &\ll_\varepsilon B^\varepsilon \sum_{v_1, v_2, y_1, y_2} \left(\frac{B^{1/4}v_2^{1/4}y_1^{1/5}}{y_2^{1/2}} + v_1^{1/4}v_2^{1/6}y_1^{1/30}B^{1/3} \right) \\ &\ll_\varepsilon B^{17/20+\varepsilon}, \end{aligned}$$

on noting that $B^{1/4}y_1^{3/20} \leq B^{1/4+3/40} \leq B^{1/3}$ and summing first over y_1 . This completes the proof of Lemma 7. $\square$

Recall the definitions (4.1), (4.2) of Y_0, Y_3 , and write $f(t) = \sqrt{1-t^4}$ as above. Our next task is to examine the sum

$$\sum_{\substack{y_0 \leq Y_0 \\ \gcd(y_0, v_1 v_2 y_1) = 1}} \frac{Y_3}{v_2 y_1^2} = \frac{B^{1/2}}{v_2^{1/2} y_1} \sum_{\substack{y_0 \leq Y_0 \\ \gcd(y_0, v_1 v_2 y_1) = 1}} f(y_0/Y_0),$$

that appears in (4.12). On performing a Möbius inversion we easily deduce that

$$\sum_{\substack{y_0 \leq Y_0 \\ \gcd(y_0, v_1 v_2 y_1) = 1}} f(y_0/Y_0) = \sum_{k_0 | v_1 v_2 y_1} \mu(k_0) \sum_{y'_0 \leq Y'_0} f(y'_0/Y'_0),$$

where $Y'_0 = Y_0/k_0$. An application of partial summation, together with the change of variables $t = u^{1/4}Y'_0$, reveals that

$$\begin{aligned} \sum_{y'_0 \leq Y'_0} f(y'_0/Y'_0) &= [Y'_0]f(1) - \int_0^{Y'_0} [t]f'\left(\frac{t}{Y'_0}\right) \frac{dt}{Y'_0} \\ &= - \int_0^1 [u^{1/4}Y'_0]f'(u^{1/4}) \frac{du}{4u^{3/4}} = cY'_0 + F_2(Y'_0), \end{aligned}$$

where

$$F_2(Y'_0) = -\frac{1}{2} \int_0^1 \{u^{1/4}Y'_0\} \frac{du}{\sqrt{1-u}},$$

and

$$c = \int_0^1 \frac{u^{1/4} du}{2\sqrt{1-u}}. \quad (4.16)$$

We easily conclude that

$$\sum_{\substack{y_0 \leq Y_0 \\ \gcd(y_0, v_1 v_2 y_1) = 1}} \frac{Y_3}{v_2 y_1^2} = c \frac{\phi(v_1 v_2 y_1)}{v_1 v_2 y_1} \frac{B^{3/4}}{v_2^{1/4} y_1^{1/2} y_2^{1/2}} + \frac{R_2(B; \mathbf{v}, y_1, y_2)}{\vartheta(\mathbf{v}, y_1, y_2)},$$

where

$$R_2(B; \mathbf{v}, y_1, y_2) = \vartheta(\mathbf{v}, y_1, y_2) \frac{B^{1/2}}{v_2^{1/2} y_1} \sum_{k_0 | v_1 v_2 y_1} \mu(k_0) F_2(Y'_0). \quad (4.17)$$

We now need to sum $R_2(B; \mathbf{v}, y_1, y_2)$ over all of the relevant values of y_2 . Let

$$Y_2 = \frac{B^{1/2}}{v_1^2 v_2^{3/2} y_1}.$$

Then the following result holds.

Lemma 8. *Let $\varepsilon > 0$ and let $B \geq 1$. Then we have*

$$\sum_{\substack{y_2 \leq Y_2 \\ \gcd(y_2, v_2 y_1) = 1}} R_2(B; \mathbf{v}, y_1, y_2) = \vartheta'(\mathbf{v}, y_1) \frac{B}{v_1^2 v_2^2 y_1^2} + O_\varepsilon \left(\frac{B^{3/4+\varepsilon}}{v_1 v_2^{5/4} y_1^{3/2}} \right),$$

where

$$\begin{aligned} \vartheta'(\mathbf{v}, y_1) &= -\frac{3}{\pi^2} \eta(v_2 y_1^2) \prod_{p | v_1 v_2} \left(1 - \frac{\chi(p)}{p} \right) \prod_{p | v_1 v_2 y_1} \left(1 + \frac{1}{p} \right)^{-1} \\ &\quad \times \sum_{k_0 | v_1 v_2 y_1} \mu(k_0) \int_0^1 \int_0^1 \left\{ \frac{u^{1/4} v_1 v_2 y_1}{k_0 v^{1/2}} \right\} \frac{du dv}{\sqrt{1-u}}. \end{aligned}$$

Proof. Now it is plain from (4.17) that

$$\sum_{\substack{y_2 \leq Y_2 \\ \gcd(y_2, v_2 y_1) = 1}} R_2(B; \mathbf{v}, y_1, y_2) = \vartheta(\mathbf{v}, y_1, 1) \frac{B^{1/2}}{v_2^{1/2} y_1} \sum_{k_0 | v_1 v_2 y_1} \mu(k_0) \sum_{y_2 \leq Y_2} \varpi(y_2) h\left(\frac{y_2}{Y_2}\right),$$

where

$$\varpi(y_2) = \begin{cases} \vartheta(\mathbf{v}, y_1, y_2) / \vartheta(\mathbf{v}, y_1, 1), & \text{if } \gcd(y_2, v_2 y_1) = 1, \\ 0, & \text{otherwise,} \end{cases}$$

and

$$h(v) = -\frac{1}{2} \int_0^1 \left\{ \frac{u^{1/4} v_1 v_2 y_1}{k_0 v^{1/2}} \right\} \frac{du}{\sqrt{1-u}}.$$

From (4.14), it follows that

$$\varpi(y_2) = \frac{\phi(y_2)}{y_2} \prod_{p | \gcd(v_1, y_2)} \left(1 - \frac{1 + \chi(p)}{p} \right)^{-1},$$

if $\gcd(y_2, v_2 y_1) = 1$. In order to estimate $\sum_{y_2 \leq Y_2} \varpi(y_2) h(y_2/Y_2)$, we must first calculate the corresponding Dirichlet series $F(s) = \sum_{n=1}^{\infty} \varpi(n) n^{-s}$. Let

$$\varepsilon_p = \left(1 - \frac{1 + \chi(p)}{p} \right)^{-1},$$

for any prime p . Then for $\Re(s) > 1$ we see that

$$F(s) = \prod_{p \nmid v_1 v_2 y_1} \left(1 + \frac{1 - 1/p}{p^s - 1} \right) \prod_{p | v_1, p \nmid v_2 y_1} \left(1 + \frac{\varepsilon_p (1 - 1/p)}{p^s - 1} \right) = \zeta(s) G(s),$$

where

$$G(s) = \prod_p \left(1 - \frac{1}{p^{s+1}} \right) \prod_{p | v_1 v_2 y_1} \left(1 + \frac{1 - 1/p}{p^s - 1} \right)^{-1} \prod_{p | v_1, p \nmid v_2 y_1} \left(1 + \frac{\varepsilon_p (1 - 1/p)}{p^s - 1} \right).$$

In particular it is clear that $F(s)/\zeta(s)$ is holomorphic for $\Re(s) > 0$. It therefore follows from a standard argument (see [2, Lemma 2] for example) that

$$\sum_{y_2 \leq t} \varpi(y_2) = G(1)t + O(G^+(1/2)t^{1/2}),$$

for any $t \geq 1$, where if $G(s) = \sum_{n=1}^{\infty} g(n)n^{-s}$ then $G^+(s) = \sum_{n=1}^{\infty} |g(n)|n^{-s}$. A simple calculation reveals that $G^+(1/2) \ll 2^{\omega(v_1 v_2 y_1)} \ll_{\varepsilon} B^{\varepsilon}$, and so an application of partial summation yields

$$\begin{aligned} \sum_{y_2 \leq Y_2} \varpi(y_2) h(y_2/Y_2) &= -\frac{G(1)}{2} \int_0^{Y_2} \int_0^1 \left\{ \frac{u^{1/4} v_1 v_2 y_1}{k_0 (s/Y_2)^{1/2}} \right\} \frac{du ds}{\sqrt{1-u}} + O_{\varepsilon}(B^{\varepsilon} Y_2^{1/2}) \\ &= -\frac{G(1)}{2} Y_2 \int_0^1 \int_0^1 \left\{ \frac{u^{1/4} v_1 v_2 y_1}{k_0 v^{1/2}} \right\} \frac{du dv}{\sqrt{1-u}} + O_{\varepsilon}(B^{\varepsilon} Y_2^{1/2}). \end{aligned}$$

Since $\vartheta(\mathbf{v}, y_1, 1) \ll 2^{\omega(v_1 y_1)} 4^{\omega(v_2)} \ll_{\varepsilon} B^{\varepsilon}$, this therefore completes the proof of Lemma 8. $\square$

While the precise value of $\vartheta'(\mathbf{v}, y_1)$ in Lemma 8 is perhaps unimportant, we will need the observation that

$$\vartheta'(\mathbf{v}, y_1) \ll 8^{\omega(v_1 v_2 y_1)} \ll_{\varepsilon} (v_1 v_2 y_1)^{\varepsilon},$$

for any $\varepsilon > 0$. We are now ready to sum $R_2(B; \mathbf{v}, y_1, y_2)$ over all $(\mathbf{v}, y_1) \in \mathbb{N}^3$ satisfying $|\mu(v_2)| = 1$ and the inequality $v_1^4 v_2^3 y_1^2 \leq B$, that follows from (4.3). Thus Lemma 8 implies that for any $\varepsilon > 0$ we have

$$\begin{aligned} \sum_{\substack{v_1, v_2, y_1 \in \mathbb{N} \\ v_1^4 v_2^3 y_1^2 \leq B}} |\mu(v_2)| R_2(B; \mathbf{v}, y_1, y_2) &= B \sum_{\substack{v_1, v_2, y_1 \in \mathbb{N} \\ v_1^4 v_2^3 y_1^2 \leq B}} \frac{|\mu(v_2)| \vartheta'(\mathbf{v}, y_1)}{v_1^2 v_2^2 y_1^2} + O_{\varepsilon}(B^{3/4+\varepsilon}) \\ &= \beta B + O_{\varepsilon}(B^{3/4+\varepsilon}), \end{aligned}$$

with

$$\beta = \sum_{v_1, v_2, y_1 \in \mathbb{N}} \frac{|\mu(v_2)| \vartheta'(\mathbf{v}, y_1)}{v_1^2 v_2^2 y_1^2}. \quad (4.18)$$

Let $(\mathbf{v}, y_1, y_2) \in \mathbb{N}^4$ satisfy (4.3), and recall the definition (4.14) of $\vartheta(\mathbf{v}, y_1, y_2)$. We define $\varphi(\mathbf{v}, y_1, y_2)$ to be zero if (3.14) fails to hold and

$$\varphi(\mathbf{v}, y_1, y_2) = \frac{\phi(v_1 v_2 y_1)}{v_1 v_2 y_1} \vartheta(\mathbf{v}, y_1, y_2) \quad (4.19)$$

otherwise. Furthermore, let

$$R(B; \mathbf{v}, y_1, y_2) = R_1(B; \mathbf{v}, y_1, y_2) + R_2(B; \mathbf{v}, y_1, y_2), \quad (4.20)$$

where $R_1(B; \mathbf{v}, y_1, y_2)$ is given by (4.13) and $R_2(B; \mathbf{v}, y_1, y_2)$ is given by (4.17). Then we have proved the following result.

Lemma 9. *Let $\varepsilon > 0$. Then for any $B \geq 1$ and any $(\mathbf{v}, y_1, y_2) \in \mathbb{N}^4$ satisfying (4.3), we have*

$$S(\mathbf{v}, y_1, y_2) = c \varphi(\mathbf{v}, y_1, y_2) \frac{B^{3/4}}{v_2^{1/4} y_1^{1/2} y_2^{1/2}} + R(B; \mathbf{v}, y_1, y_2),$$

where c is given by (4.16), $\varphi(\mathbf{v}, y_1, y_2)$ is given by (4.19), and $R(B; \mathbf{v}, y_1, y_2)$ is given by (4.20) and satisfies

$$\sum_{\substack{v_1, v_2, y_1, y_2 \in \mathbb{N} \\ (3.14), (4.3) \text{ hold}}} R(B; \mathbf{v}, y_1, y_2) = \beta B + O_{\varepsilon}(B^{17/20+\varepsilon}),$$

where β is given by (4.18).

4.3. Summation over the remaining variables. In this section we complete our preliminary estimate for $N_{U,H}(B)$, for which we first recall the definition (3.1) of the counting function $N(Q_1, Q_2; B)$. Our task is to sum the main term in Lemma 9 over all $\mathbf{v}, y_1, y_2$ satisfying (3.14) and (4.3). Define the arithmetic function

$$\Delta(n) = \sum_{\substack{\mathbf{v}, y_1, y_2 \in \mathbb{N} \\ v_1^4 v_2^3 y_1^2 y_2^2 = n}} \frac{\varphi(\mathbf{v}, y_1, y_2)}{v_2^{1/4} y_1^{1/2} y_2^{1/2}}, \quad (4.21)$$

for any $n \in \mathbb{N}$. Then it follows from Lemma 9 that

$$\begin{aligned} N(Q_1, Q_2; B) &= cB^{3/4} \sum_{\substack{v_1, v_2, y_1, y_2 \in \mathbb{N} \\ v_1^4 v_2^3 y_1^2 y_2^2 \leq B}} \frac{\varphi(v_1, v_2, y_1, y_2)}{v_2^{1/4} y_1^{1/2} y_2^{1/2}} + \beta B + O_\varepsilon(B^{17/20+\varepsilon}) \\ &= cB^{3/4} \sum_{n \leq B} \Delta(n) + \beta B + O_\varepsilon(B^{17/20+\varepsilon}), \end{aligned}$$

for any $\varepsilon > 0$. On inserting this estimate into Lemma 2 we therefore obtain the following result.

Lemma 10. *Let $\varepsilon > 0$. Then for any $B \geq 1$ we have*

$$N_{U,H}(B) = 4cB^{3/4} \sum_{n \leq B} \Delta(n) + \left(\frac{12}{\pi^2} + 4\beta\right)B + O_\varepsilon(B^{17/20+\varepsilon}),$$

where c is given by (4.16), $\Delta(n)$ is given by (4.21) and β is given by (4.18).

5. THE HEIGHT ZETA FUNCTION

In this section we complete the proof of Theorem 1. Recall the definition (4.21) of $\Delta(n)$, which has corresponding Dirichlet series

$$D(s) = \sum_{n=1}^{\infty} \frac{\Delta(n)}{n^s} = \sum_{v_1, v_2, y_1, y_2 \in \mathbb{N}} \frac{\varphi(v_1, v_2, y_1, y_2)}{v_1^{4s} v_2^{3s+1/4} y_1^{2s+1/2} y_2^{2s+1/2}},$$

say, where $\varphi(v_1, v_2, y_1, y_2)$ is given by (4.19). For $\Re(s) > 1$ the height zeta function is given by

$$Z_{U,H}(s) = \sum_{x \in U(\mathbb{Q})} \frac{1}{H(x)^s} = s \int_1^\infty t^{-s-1} N_{U,H}(t) dt.$$

Thus it follows from Lemma 10 that $Z_{U,H}(s) = Z_1(s) + Z_2(s)$, where

$$\begin{aligned} Z_1(s) &= 4cs \int_1^\infty t^{-s-1/4} \sum_{n \leq t} \Delta(n) dt = \frac{16cs}{4s-3} D(s-3/4), \\ Z_2(s) &= \frac{12/\pi^2 + 4\beta}{s-1} + G_2(s), \end{aligned}$$

and

$$G_2(s) = s \int_1^\infty t^{-s-1} R(t) dt$$

for some function $R(t)$ such that $R(t) \ll_\varepsilon t^{17/20+\varepsilon}$ for any $\varepsilon > 0$. But then it follows that $G_2(s)$ is holomorphic on the half-plane $\{s \in \mathbb{C} : \Re(s) \geq 17/20 + \varepsilon\}$, and is easily seen to satisfy the inequality

$$G_2(s) \ll_\varepsilon (1 + |\Im(s)|)^{20 \max\{1-\Re(s), 0\} + 3 + \varepsilon}.$$

on this domain, via the Phragmén-Lindelöf Theorem.

It remains to analyse the function $Z_1(s)$. The multiple sum $D(s)$ can be written as an Eulerian product $\prod_p D_p(s)$, say. When p is odd a routine calculation reveals that

$$D_p(s+1/4) = 1 + \frac{(1-1/p)(2+\chi(p))}{p^{1+2s}-1} \left(1 + \frac{1-1/p}{p^{1+4s}-1}\right) \\ + \frac{(1-1/p)(1-(1+\chi(p))/p)}{p^{1+4s}-1} + \frac{(1-1/p)^2(1+\chi(p))}{p^{1+3s}(1-p^{-1-4s})(1-p^{-1-2s})}.$$

Similarly, in the case $p=2$ we have

$$D_2(s+1/4) = 1 + \frac{1}{2^{1+2s}-1} \left(\frac{1}{2} + \frac{1}{4(2^{1+4s}-1)}\right) + \frac{1}{4(2^{1+4s}-1)} \left(1 + \frac{1}{2^{3s}}\right) + \frac{1}{2^{2+3s}}.$$

Recall the definition (1.4) of $E_1(s+1)$. We have therefore shown that there exists a function H , which is analytic on the half plane $\Re(s) > 1/8$, such that

$$D(s+1/4) = E_1(s+1)H(s). \quad (5.1)$$

Let $H_p(s)$ denote the Eulerian factor of $H(s)$ at the prime p . We examine the analytic properties of $H(s)$ by estimating the size of $H_p(s)$. In doing so we restrict ourselves to the domain $\mathcal{D} = \{s \in \mathbb{C} : \Re(s) \geq -1/4 + \varepsilon\}$. When p is odd we plainly have

$$D_p(s+1/4)(1-p^{-1-4s})(1-p^{-1-2s}) = 1 + \frac{1+\chi(p)}{p^{1+2s}} + \frac{1+\chi(p)}{p^{1+3s}} + O\left(\frac{1}{p^{1+\varepsilon}}\right).$$

It therefore follows from (5.1) that

$$\frac{H_p(s)}{(1-p^{-1-3s})(1-\chi(p)p^{-1-3s})} = 1 + \frac{1+\chi(p)}{p^{1+3s}} - \frac{2(1+\chi(p))}{p^{2+5s}} + O\left(\frac{1}{p^{1+\varepsilon}}\right),$$

whence

$$H_p(s) = 1 - \frac{2+\chi(p)}{p^{2+6s}} - \frac{2(1+\chi(p))}{p^{2+5s}} + \frac{1+\chi(p)}{p^{3+9s}} + O\left(\frac{1}{p^{1+\varepsilon}}\right) \\ = E_{2,p}(s+1) \left(1 + O\left(\frac{1}{p^{1+\varepsilon}}\right)\right)$$

on $\mathcal{D}$. Here $E_{2,p}(s+1)$ is the Eulerian factor of $E_2(s+1)$, as given by (1.5). Define the function

$$G_1(s) = \frac{16cs}{4s-3} \frac{D(s-3/4)}{E_1(s)E_2(s)} = \frac{16cs}{4s-3} \frac{H(s-1)}{E_2(s)}. \quad (5.2)$$

Then it follows from our work that $G_1(s)$ is analytic and bounded on the half-plane $\{s \in \mathbb{C} : \Re(s) \geq 3/4 + \varepsilon\}$. In order to complete the proof of Theorem 1 we simply note from (5.1) that

$$H(0) = \frac{5}{2^5} \prod_{p>2} \left(1 - \frac{1}{p}\right)^4 \left(1 - \frac{\chi(p)}{p}\right)^2 \left(1 + \frac{4+2\chi(p)}{p} + \frac{1}{p^2}\right) = \tau, \quad (5.3)$$

by (2.6), whence

$$G_1(1) = 16c \frac{H(0)}{E_2(1)} \neq 0.$$

6. DEDUCTION OF THEOREM 2

This section is virtually identical to the corresponding argument in [2, §7], and so we will be brief. Let $\varepsilon > 0$ and let $T \in [1, B]$. Then it follows from an application of Theorem 1, Lemma 10 and Perron's formula that

$$N_{U,H}(B) - \left(\frac{12}{\pi^2} + 4\beta\right)B = \frac{1}{2\pi i} \int_{1+\varepsilon-iT}^{1+\varepsilon+iT} E_1(s)E_2(s)G_1(s) \frac{B^s}{s} ds + O_\varepsilon\left(\frac{B^{1+\varepsilon}}{T} + B^{17/20+\varepsilon}\right). \quad (6.1)$$

We apply Cauchy's residue theorem to the rectangular contour $\mathcal{C}$ joining the points $\kappa - iT$, $\kappa + iT$, $1 + \varepsilon + iT$ and $1 + \varepsilon - iT$, for any $\kappa \in [29/32, 1)$. On expanding the product of zeta functions about $s = 1$ it follows from (1.4) that

$$E_1(s) = \frac{1}{48(s-1)^4} + O\left(\frac{1}{(s-1)^3}\right),$$

from which it is easy to deduce from (5.2) and (5.3) that

$$\text{Res}_{s=1} \left\{ E_1(s)E_2(s)G_1(s) \frac{B^s}{s} \right\} = \frac{c}{3! \times 4} L(1, \chi)^2 \tau B Q(\log B),$$

for some monic polynomial Q of degree 3.

Define the difference

$$E(B) = N_{U,H}(B) - \frac{c}{3! \times 4} L(1, \chi)^2 \tau B Q(\log B) - \left(\frac{12}{\pi^2} + 4\beta\right)B.$$

Then, in view of (6.1) and the fact that the function $E_2(s)G_1(s)$ is holomorphic and bounded for $\Re(s) > 5/6$, we deduce that

$$E(B) \ll_\varepsilon \frac{B^{1+\varepsilon}}{T} + B^{17/20+\varepsilon} + \left(\int_{\kappa-iT}^{\kappa+iT} + \int_{\kappa-iT}^{1+\varepsilon-iT} + \int_{1+\varepsilon+iT}^{\kappa+iT} \right) \left| E_1(s) \frac{B^s}{s} \right| ds, \quad (6.2)$$

for any $\kappa \in [29/32, 1)$ and any $T \in [1, B]$. We begin by estimating the contribution from the horizontal contours. Recall the well-known convexity bounds

$$\zeta(\sigma + it) \ll_\varepsilon |t|^{(1-\sigma)/3+\varepsilon}, \quad L(\sigma + it, \chi) \ll_\varepsilon |t|^{1-\sigma+\varepsilon},$$

that are valid for any $\sigma \in [1/2, 1]$ and $|t| \geq 1$. Then it follows that

$$E_1(\sigma + it) \ll_\varepsilon |t|^{26(1-\sigma)/3+\varepsilon}$$

for any $\sigma \in [29/32, 1]$ and $|t| \geq 1$. This estimate allows us to deduce that

$$\left(\int_{\kappa-iT}^{1+\varepsilon-iT} + \int_{1+\varepsilon+iT}^{\kappa+iT} \right) \left| E_1(s) \frac{B^s}{s} \right| ds \ll_\varepsilon \frac{B^{1+\varepsilon} T^\varepsilon}{T} + B^\kappa T^\varepsilon. \quad (6.3)$$

We now turn to the size of the integral

$$\int_{\kappa-iT}^{\kappa+iT} \left| E_1(s) \frac{B^s}{s} \right| ds \ll B^\kappa \int_{-T}^T \frac{|E_1(\kappa + it)|}{1 + |t|} dt = B^\kappa I(T), \quad (6.4)$$

say. For given $0 < U \ll T$, we will estimate the contribution to $I(T)$ from each integral

$$\int_U^{2U} \frac{|E_1(\kappa + it)|}{1 + |t|} dt \ll \frac{1}{U} \int_U^{2U} |E_1(\kappa + it)| dt = \frac{J(U)}{U},$$

say. For any $\varepsilon > 0$ it follows from a result due to Heath-Brown [8] that

$$\int_U^{2U} |\zeta(\sigma + it)|^8 dt \ll_\varepsilon U^{1+\varepsilon}, \quad (6.5)$$

for any $\sigma \in [5/8, 1]$, and any $U \geq 1$. Moreover we will need the upper bound

$$\int_U^{2U} |L(\sigma + it, \chi)|^4 dt \ll_\varepsilon U^{1+\varepsilon}, \quad (6.6)$$

that is valid for any $\sigma \in [1/2, 1]$ and any $U \geq 1$. This follows on combining standard convexity estimates [16, §7.8] with work of Montgomery [13, Theorem 10.1]. Returning to our estimate for $J(U)$, for fixed $0 < U \ll T$ and any $\kappa \in [29/32, 1]$, we apply Hölder's inequality to deduce that

$$J(U) \leq J_1^{1/8} J_2^{1/8} J_3^{1/4} J_4^{1/4} J_5^{1/4},$$

where

$$\begin{aligned} J_1 &= \int_U^{2U} |\zeta(4\kappa - 3 + 4it)|^8 dt, & J_2 &= \int_U^{2U} |\zeta(3\kappa - 2 + 3it)|^8 dt, \\ J_3 &= \int_U^{2U} |\zeta(2\kappa - 1 + 2it)|^8 dt, & J_4 &= \int_U^{2U} |L(3\kappa - 2 + 3it, \chi)|^4 dt, \\ J_5 &= \int_U^{2U} |L(2\kappa - 1 + 2it, \chi)|^4 dt. \end{aligned}$$

It therefore follows from (6.5) and (6.6) that $J(U) \ll_\varepsilon U^{1+\varepsilon}$, on re-defining the choice of ε . Summing over dyadic intervals for $0 < U \ll T$ we obtain

$$\int_0^T \frac{|E_1(\kappa + it)|}{1 + |t|} dt \ll_\varepsilon T^\varepsilon.$$

Since we obtain the same estimate for the integral over the interval $[-T, 0]$, we deduce that $I(T) \ll_\varepsilon T^\varepsilon$. We may insert this estimate into (6.4), and then combine it with (6.3) in (6.2), in order to conclude that

$$E(B) \ll_\varepsilon \frac{B^{1+\varepsilon}}{T} + B^{\kappa+\varepsilon},$$

for any $T \in [1, B]$. We therefore complete the proof of Theorem 2 by taking $T = B$.

REFERENCES

- [1] V.V. Batyrev and Y. Tschinkel, Manin's conjecture for toric varieties. *J. Alg. Geom.* **7** (1998), 15–53.
- [2] R. de la Bretèche and T.D. Browning, On Manin's conjecture for singular del Pezzo surfaces of degree four, I. *Michigan Math. J.* **55** (2007), 51–80.
- [3] T.D. Browning, An overview of Manin's conjecture for del Pezzo surfaces. *Proceedings of the Gauss–Dirichlet conference (Göttingen)*, to appear.
- [4] A. Chambert-Loir and Y. Tschinkel, On the distribution of points of bounded height on equivariant compactifications of vector groups. *Invent. Math.* **148** (2002), 421–452.
- [5] D.F. Coray and M.A. Tsfasman, Arithmetic on singular Del Pezzo surfaces. *Proc. London Math. Soc.* **57** (1988), 25–87.
- [6] U. Derenthal and Y. Tschinkel, Universal torsors over del Pezzo surfaces and rational points. *Equidistribution in Number Theory: An Introduction (Montreal, 2005)*, 169–196, NATO Science Series II: Math, Physics and Chemistry **237**, Springer, 2006.
- [7] J. Franke, Y.I. Manin and Y. Tschinkel, Rational points of bounded height on Fano varieties. *Invent. Math.* **95** (1989), 421–435.
- [8] D.R. Heath-Brown, Mean values of the zeta-function and divisor problems. *Recent progress in analytic number theory*, Vol. I, 115–119, Academic Press, 1981.
- [9] D.R. Heath-Brown, Weyl's inequality, Hua's inequality, and Waring's problem. *J. London Math. Soc.* **38** (1988), no. 2, 216–230.
- [10] B. Hassett and Y. Tschinkel, Universal torsors and Cox rings. *Arithmetic of higher-dimensional algebraic varieties (Palo Alto, 2002)*, 149–173, Progr. Math. **226**, Birkhäuser, 2004.
- [11] W.V.D. Hodge and D. Pedoe, *Methods of algebraic geometry*. Vol. 2, Cambridge University Press, 1952.

- [12] J. Lipman, Rational singularities, with applications to algebraic surfaces and unique factorization. *Inst. Hautes Études Sci. Publ. Math.* **36** (1969), 195–279.
- [13] H.L. Montgomery, *Topics in multiplicative number theory*. Springer Lecture Notes 227, Springer, 1971.
- [14] E. Peyre, Hauteurs et mesures de Tamagawa sur les variétés de Fano. *Duke Math. J.* **79** (1995), 101–218.
- [15] P. Swinnerton-Dyer, Counting points on cubic surfaces, II. *Geometric methods in algebra and number theory*, 303–310, Progr. Math. **235**, Birkhäuser, 2005.
- [16] E.C. Titchmarsh, *The theory of the Riemann zeta-function*. 2nd ed., edited by D.R. Heath-Brown, Oxford University Press, 1986.
- [17] J.D. Vaaler, Some extremal functions in Fourier analysis. *Bull. Amer. Math. Soc.* **12** (1985), no. 2, 183–216.

INSTITUT DE MATHÉMATIQUES DE JUSSIEU, UNIVERSITÉ PARIS 7 DENIS DIDEROT, CASE POSTALE
7012, 2, PLACE JUSSIEU, F-75251 PARIS CEDEX 05
E-mail address: `breteche@math.jussieu.fr`

SCHOOL OF MATHEMATICS, UNIVERSITY OF BRISTOL, BRISTOL BS8 1TW
E-mail address: `t.d.browning@bristol.ac.uk`